

## **BioCLIA Autoimmune Calibrator Set, Ro52**

Cat.No.	Kit Size
MY00206	2 X 1 mL
MY00257	4 X 1 mL

## **INTENDED USE**

The BioCLIA Autoimmune Calibrator Set, Ribosomal P is intended for the calibration of the BioCLIA Ribosomal P performed on the BioCLIA 6500 and BioCLIA 500.

For professional in vitro diagnostic use only.

## **SUMMARY AND EXPLANATION**

Anti-nuclear antibody (ANA) is a class of auto-antibodies with different binding affinities specific to different nuclear antigens. Generally, ANAs include extractable nuclear antigen (ENA) antibodiesand un-extractable nuclear antigen antibodie.¹ Determination of ANA has significant correlation to the diagnosis of Sharp syndrome (MCTD), systemic lupus erythematosus (SLE), sjogren's syndrome, progressive systemic sclerosis, polymyositis/dermatomyositis, overlap syndrome, andlimited types of progressive systemic sclerosis (CREST syndrome).

Ro52 is a 52 kDa protein detected in autoimmune diseases patients' sera include sjogren's syndrome, systemic lupus 7erythematosus (SLE), dermatomyositis, *etc.* It belongs to the TRIM proteinfamily(TRIM21) with different functional structure domain like RING finger, B-box or curly spiral. <sup>2,3</sup> The effect of Ro52 on immune defense is increasingly concerned, such as the interaction between Ro52 and IgG molecules, or how the Ro52 being an E3 ubiquitin ligase and act on signal molecules. So far, the pathogenic mechanism of Ro52in autoimmune diseases is not clear.Ro52 proteins are confirmed to be associated with SS-A/Ro complex function. <sup>4</sup>

Due to the appearance of anti-Ro52 antibodies in a variety of autoimmune diseases, it should not been taken as a specific indicator of SLE or sjogren's syndrome but should combine with other clinical findings.

# MATERIALS SUPPLIED

• Ro52 Calibrator 1 Barcode labeled tubes with buffer containing human antibodies to Ro52 in stabilizers and preservatives. Ready to use, 1 mL.

CAL 1

Preservatives: 0.0015% < Proclin 300 < 0.6%.

• Ro52 Calibrator 2 Barcode labeled tubes with buffer containing human antibodies to Ro52 in stabilizers and preservatives. Ready to use, 1 mL.

CAL 2

Preservatives: 0.0015% < Proclin 300 < 0.6%.

The Calibrator Code contains calibrators' information is provided in each kit.

# WARNINGS AND PRECAUTIONS

• For professional in vitro diagnostic use only.

- Do not use any calibrators beyond their expiration dates.
- Do not mix calibrators from different lots unless specified.
- Instructions must be carefully followed for using and storing of calibrators. Any modification in procedure may interfere with the results.
- Calibrators and contaminated vials must be handled strictly following safety guidelines or rules of biological hazards to ensure the users' and environmental safety.
- Any serious incident that has occurred in relation to the device shall be reported to the manufacturer and the competent authority of the Member State in which the user and/or the patient is established!

#### **Precautions:**



Human serum is added in the calibrators.

The human derived material in this product was tested by FDA approved methods and found nonreactive for Hepatitis B Surface Antigen (HBsAg), Anti-HCV and HIV 1/2 antibodies. Handle as if potentially infectious. <sup>7</sup> Avoid contacting with skin and eyes. Do not empty into drains. Wear suitable protective gloves and clothing.



- Proclin 300 is added in the calibrators at concentration between 0.0015% 0.6%.
- Calibrators contain chemical and biological components. Avoid ingesting or splashing onto skin and mucous membrane. If direct contact with calibrators happens, rinse immediately the contact surface with plenty of water and see a doctor if necessary.

# STORAGE CONDITIONS

- Store the kit at 2-8 °C.
- The shelf life of the unopened kit is 12 months from day of production.
- Vial opened calibrators could be used for 28 successive days, exposure no more than 2 hours each time when kept uncapped and is good for up to 20 calibrations, after which the reagent must be discarded.
- Avoid repeated freezing and thawing.

# **ASSAY PROCEDURE**

Detailed information about operating the BioCLIA instruments can be

1



taken from the Instrument User's Manual.

Note that, it is important to perform all routine maintenance procedures for optimal performance.

## **Assay Calibration**

The BioCLIA Autoimmune Reagent Kit utilizes a predefined lot specific Master Curve which is uploaded into the instrument via the barcode provided in the main reagent kit. The Calibrator Code contains calibrator information is then scanned. Based on the results of running two calibrators, the instrument specific Working Calibration Curve is generated and is used to calculate the concentration from the RLU obtained for each patient.

For each new lot of reagent, please calibrate prior to the first time use, and every 28 days thereafter. The software will not allow the lot to be used if the above requirements are not meet.

# **Programming and Running samples**

- Put the kit into the corresponding position of the reagent chamber of the fully automatic chemiluminescence analyzer. The information of the kit can be uploaded into the instrument system through the scanning of reagent barcode, and can also be set through the supporting software of the instrument.
- The information of calibrator / quality control is identified by scanning the calibrator / control barcodes, and the position of calibrator / quality control is assigned in the instrument system.
- The sample to be tested is placed on the instrument sample rack chamber, and the corresponding test information is edited through the instrument supporting software.
- 4. Start the operation procedure, and all calibrator / quality control / sample processing steps will be automatically executed.

## **TRACEABILITY**

The reported values were determined over multiple runs on the BioCLIA 6500 and BioCLIA 500 using specific lots of reagents against an in-house standard. DGP IgA results are reported in RU/mL which is interpreted from relative light unit (RLU). Method comparison test showed good sensitivity and specificity of tested assay.

## **LIMITATIONS**

- The calibrators are designed for calibration of the same lot of BioCLIA Autoimmune Reagent Kit.
- The calibrators can be kept uncapped onboard the instrument up to 2 hours for each time of usage. And a total up to 20 calibrations are suggested, for any longer period of time, the reagent should be discarded, otherwise may result in improper calibration of the assay and which can give improper results.

## **SYMBOLS**

CAL 1	Calibrator 1	
CAL 2	Calibrator 2	

REF	Catalog Number	$\square$	Use-by date
IVD	In Vitro diagnostic medical device	LOT	Lot Number
+2°C	Store between +2°C and +8°C	(i	Consult Instruction for Use
***	Manufacturer	EC REP	Authorized Representative in the European Community
(€	CE Marking	\subseteq	Contains Sufficient for <n>Tests</n>
₩	Biological Risk	1>	GHS07 Warning

#### REFERENCE

- 1. Tan EM. Autoantibodies to nuclear antigens (ANA): their immunobiology and medicine. Advances in Immunology 1982;33:167-240.
- 2. Frank M, Itoh K, Fujisaku A, Pontarotti P, Mattei M, Neas B. The mapping of the human 52-kD Ro/SSA autoantigen gene to human chromosome 11, and its polymorphisms. American journal of human genetics 1993:52:183.
- 3. James LC, Keeble AH, Khan Z, Rhodes DA, Trowsdale J. Structural basis for PRYSPRY-mediated tripartite motif (TRIM) protein function. Proceedings of the National Academy of Sciences 2007;104:6200-05.
- 4. Mallery DL, McEwan WA, Bidgood SR, Towers GJ, Johnson CM, James LC. Antibodies mediate intracellular immunity through tripartite motif-containing 21 (TRIM21). Proceedings of the National Academy of Sciences 2010;107:19985-90.
- Richmond JY, Mckinney RW. Biosafety in microbiological and biomedical laboratories: U.S.GPO.1999.



HOB Biotech Group Corp., Ltd.

C6 Building, No. 218 Xinghu Road, Suzhou Industrial Park, Suzhou, Jiangsu, 215123, China

## **CONTACT INFORMATION:**

TEL (+86)512-69561996 Fax (+86)512-62956652

WEBSITE: www.hob-biotech.com

**CUSTOMER SERVICE:** HOB Biotech Group Co., Ltd **CUSTOMER SERVICE:** TEL (+86)4008601202

EC REP

**EUROPE REPRESENTATIVE:** Emergo Europe

ADDRESS/LOCATION:

Prinsessegracht 20, 2514 AP The Hague, The Netherlands



## The eIFU is available on Website:

http://en.hob-biotech.com/usercenter/login.aspx

## **TECHNICAL ASSISTANCE**

For technical assistance, contact your National Distributor.

Date of issue: 17<sup>th</sup> March 2019

Date of revision: 20<sup>th</sup> December 2021 Change Control Number: CN21129E

Version: A/1 (EN)