

***Fluorescein-Labeled  
Affinity Purified Antibody  
To Monkey IgG ( $\gamma$ )***

*Produced in Goat*

**Catalog No.   Size**  
**072-11-021   1.0 mg**



**DESCRIPTION**

Affinity purified antibody isolated from a pool of serum from goats immunized with purified Old World monkey IgG was labeled with fluorescein isothiocyanate (FITC) using a modified method adapted from The and Felthamp (1).

**FORM/STORAGE**

Lyophilized. Store at 2-8°C until rehydrated. Stable lyophilized for a minimum of 1 year when stored at 2-8°C.

**STABILIZER AND PRESERVATIVE**

Bovine serum albumin (BSA) is added as a protein stabilizer. A proprietary anti-bacterial agent is added to preserve the product. Non-sterile.

**ANTIBODY CONCENTRATION**

The concentration of affinity purified antibody is 1 mg as determined by UV absorbance at 280 nm.

**E/P RATIO**

Fluorescein/antibody protein ratio = 2-5:1

**SPECIFICITY/CROSS REACTIVITY**

Tested by immunoelectrophoresis techniques as applicable. This product reacts specifically with Old World monkey IgG. Product has been adsorbed with purified monkey IgA and IgM. Reactivity to IgG subclasses has not been tested. Antibodies to monkey IgG may cross-react with immunoglobulins of other mammalian species if common binding sites are shared.

**REHYDRATION AND STORAGE**

**Note:** Rehydration of antibodies in TBS or buffers other than those listed here is not recommended.

**Procedure A: 50% Glycerol**

At a working dilution, the level of glycerol is too small to affect most assays. The use of glycerol is not recommended when the conjugate is used in live cell work.

**Rehydration:** Add 0.5 mL reagent quality water to the product vial. Rotate the vial until the lyophilized pellet is totally dissolved. Add 0.5 mL glycerol to the product vial. Pipette up and down several times to ensure proper mixing. Dilute to desired concentration with TBS or other buffer.

**Storage:** This product may be stored at 2-8°C for up to 1 week; thereafter, it should be stored at -20°C.

**Procedure B: H<sub>2</sub>O**

**Rehydration:** Rehydrate with 1 mL of reagent quality water. Rotate the vial until the lyophilized pellet is totally dissolved. Dilute to desired concentration with TBS or other buffer.

**Storage:** This product may be stored for up to 1 week refrigerated; thereafter, it should be stored at -20°C.

**SUGGESTED WORKING DILUTIONS**

Optimal working concentrations should be determined experimentally. Prepare working dilution in TBS or other buffer such as BSA or Milk Diluent/Blocking Solution (See RELATED PRODUCTS). These buffers not recommended for long term storage. Suggested starting dilutions are as follows.

**Note:** In many cases, the antibody may be diluted further than indicated.

**Immunomicroscopy:**

1:100 – 1:1000                      5.0 µg/ml - 2.0 µg/ml

**Flow Cytometry:**

1:250 – 1:500                      20.0 µg/ml - 5.0 µg/ml

**Fluorescein is excited at 495 nm and emits at 528 nm.**

## PRODUCT SAFETY AND HANDLING

This product is considered non-hazardous as defined by The Hazard Communication Standard (29 CFR 1910.1200). Avoid contact with skin and eyes. In case of contact or spillage, clean with copious amounts of water. Disposal via sanitary sewer.

## REFERENCES

1. The and Felthamp, (1970). *Immunology*, 18:865.

## RELATED PRODUCTS

BSA Diluent/Blocking Solution	Cat. No. 50-61-00
Milk Diluent/Blocking Solution	Cat. No. 50-82-01
Normal Goat Serum	Cat. No. 71-00-27
Wash Solution Concentrate	Cat. No. 50-63-00
Propidium Iodide	Cat. No. 71-04-01

See the KPL catalog for a wide selection of antibodies, substrates, protein and nucleic acid detection kits, and immunohistochemistry reagents.

The product listed herein is for research use only and is not intended

for use in human or clinical diagnosis. Nothing disclosed herein is to be construed as a recommendation to use this product in violation of any patents. The information presented above is believed to be accurate. However, said information and product are offered without warranty or guarantee since the ultimate conditions of use and the variability of the materials treated are beyond our control. We cannot be responsible for patent infringements or other violations that may occur with the use of this product. No claims beyond replacement of unacceptable material or refund of purchase price shall be allowed.

L-620-03