


**ReserveAP™ Phosphatase-Labeled
Antibody To Rabbit IgG (H+L)**
(Human Serum Adsorbed)
Produced in Goat
Catalog No. Size
0751-1516 1.0 mg



DESCRIPTION

Affinity purified antibody isolated from a pool of serum from goats immunized with purified rabbit IgG was labeled with phosphatase using a modified glutaraldehyde procedure (1).

FORM/STORAGE

Lyophilized. Store at 2-8°C until rehydrated. Stable for a minimum of 1 year when stored at 2-8°C.

STABILIZER AND PRESERVATIVE

Goat serum and/or bovine serum albumin (BSA) are added as a protein stabilizer. No preservatives are added. Additional biological protection may be provided with 0.1% sodium azide. Non-sterile.

ANTIBODY CONCENTRATION

The concentration of affinity purified antibody is 1.0 mg as determined by UV absorbance at 280 nm.

E/P RATIO

Molar enzyme/antibody protein ratio is from 1.9:1 to 2.9:1.

SPECIFICITY/CROSS REACTIVITY

Tested by gel diffusion and ELISA techniques as applicable. This product reacts specifically with rabbit IgG and may recognize other immunoglobulin types that have light chains in common with IgG. Reactivity to IgG subclasses has not been tested. Antibodies to rabbit IgG may cross-react with immunoglobulins of other mammalian species if common binding sites are shared. Cross-reactivity with human serum has been minimized with affinity procedures.

REHYDRATION AND STORAGE

Note: Rehydration of antibodies in buffers other than those listed here is not recommended.

Procedure A: 50% Glycerol

At a working dilution, the level of glycerol is too small to affect most assays. The use of glycerol is not recommended when the conjugate is used in live cell work.

Rehydration: Add 0.5 mL reagent quality water to the product vial. Rotate the vial until the lyophilized pellet is totally dissolved. Add 0.5 mL glycerol to the product vial. Pipette up and down several times to ensure proper mixing.

Use: Prior to use, dilute to desired concentration with TBS or other buffer. The diluted conjugate should be used immediately.

Storage: This product may be stored either refrigerated or frozen as desired. Stable for a minimum of 1 year. **This is the best way to rehydrate a conjugate for long-term storage.**

Procedure B: H₂O

Rehydration: Rehydrate with 1 mL of reagent quality water. Rotate the vial until the lyophilized pellet is totally dissolved.

Use: Prior to use, dilute to desired concentration with TBS or other buffer. The diluted conjugate should be used immediately.

Storage: This product may be stored for up to 1 week refrigerated; thereafter, it should be stored frozen. Stable for a minimum of 1 year at -20°C.

SUGGESTED WORKING DILUTIONS

Optimal working concentrations should be determined experimentally. Prepare working dilution in TBS or other buffer such as AP Conjugate Stabilizer, BSA or Milk Diluent/Blocking Solution (See RELATED PRODUCTS). These buffers are not recommended for long-term storage, with the exception of AP Conjugate Stabilizer (See RELATED PRODUCTS). Suggested starting dilutions are as follows. In many cases, the antibody may be diluted further than indicated:

Microwell:

1:2000 - 1:10000 0.5 µg/mL - 0.1 µg/mL

Blotting:

1:10000 - 1:40000 0.1 µg/mL - 0.025 µg/mL

Histo/Cytochemical Procedures:

1:400-1:1000 2.5 µg/mL - 1.0 µg/mL

ReserveAPTM Phosphatase-Labeled Antibody To Rabbit IgG (H+L)

Catalog No. Size
0751-1516 1.0 mg

PRODUCT SAFETY AND HANDLING

This product is considered non-hazardous as defined by The Hazard Communication Standard (29 CFR 1910.1200). Avoid contact with skin and eyes. In case of contact or spillage, clean with copious amounts of water. Disposal via sanitary sewer.

REFERENCES

1. Voller, A., et. al. (1976). A Manual of Clinical Immunology. American Society for Microbiology, 509-510.

RELATED PRODUCTS

AP Conjugate Stabilizer (200 mL)	Cat. No. 55-15-00
BSA Diluent/Blocking Solution	Cat. No. 50-61-00
Milk Diluent/Blocking Solution	Cat. No. 50-82-01
Coating Solution	Cat. No. 50-84-00
Wash Solution	Cat. No. 50-63-00

Suggested KPL Phosphatase Substrates

Microwell:

BluePhos [®]	Cat. No. 50-88-00
pNPP	Cat. No. 50-80-00
RedPhos [™] Microwell	Cat. No. 50-87-20

Immunoblotting:

BCIP/NBT	Cat. No. 50-81-07
RedPhos [™] Membrane	Cat. No. 50-81-30
PhosphaGLO [™]	Cat. No. 55-60-04
PhosphaGLO Reserve [™]	Cat. No. 55-60-02

Immunohistochemistry:

HistoMark [®] RED	Cat. No. 55-69-00
HistoMark [®] BLUE	Cat. No. 55-70-00

See the KPL catalog for a wide selection of antibodies, substrates, protein and nucleic acid detection kits, and immunohistochemistry reagents.

The product listed herein is for research use only and is not intended for use in human or clinical diagnosis. Nothing disclosed herein is to be construed as a recommendation to use this product in violation of any patents. The information presented above is believed to be accurate. However, said information and product are offered without warranty or guarantee since the ultimate conditions of use and the variability of the materials treated are beyond our control. We cannot be responsible for patent infringements or other violations that may occur with the use of this product. No claims beyond replacement of unacceptable material or refund of purchase price shall be allowed. All claims regarding product performance must be made within 30 days following date of delivery.

Limited Use License

The purchase of this product conveys to the buyer the non-transferable right to use the product in research conducted by the buyer. The buyer cannot sell or otherwise transfer this product or materials made by use of this product to a third party or otherwise use this product or materials made with this product for Commercial Purposes without written approval of KPL, Inc. For additional information, please visit www.kpl.com or the KPL catalog. To obtain a license or approval to use this product for purposes other than those permitted above, contact Director of Sales at (301) 948-7755.