



ImmunoDNA Digestor





Intended Use

For Research Use Only.

Summary and Explanation

ImmunoDNA Digestor is used for the enzymatic digestion of paraffin-embedded, formalin-fixed tissues prior to immunohistochemical (IHC) or in situ hybridization (ISH) procedures.

The enzymatic digestion of formalin-fixed tissues improves accessibility of antibodies and DNA probes to target sites within tissue sections. In the case of IHC, proteolytic digestion exposes certain epitopes which have been masked during fixation. In ISH procedures, accessibility of tissue nucleic acid sequences is enhanced, allowing better probe penetration and hybridization.

Presentation

ImmunoDNA Digestor is provided in liquid form ready-to-use. It contains optimized Proteinase K, a buffer solution, detergents, and a sodium azide anti-microbial.

Catalog No.	Concentration	Volume
BSB-0108-RUO	Ready-to-use	15 mL
BSB-0109-RUO	Ready-to-use	50 mL
BSB-0110-RUO	Ready-to-use	100 mL
BSB-0111-RUO	Ready-to-use	200 mL
BSB-0112-RUO	Ready-to-use	1000 mL

Storage Store at 2-8°C

Stability

This product is stable up to the expiration date on the product label.

Do not use after expiration date listed on package label. Temperature fluctuations should be avoided. Store appropriately when not in use. Adhere to all local laws when disposing of this product.

Precautions

- 1. For professional users only. Results should be interpreted by a qualified medical professional.
- 2. This product contains <0.1% sodium azide (NaN₃) as a preservative. Ensure proper handling procedures are used with this reagent.
- 3. Always wear personal protective equipment such as laboratory coat, goggles and gloves when handling reagents.
- 4. Dispose of unused solution with copious amount of water.
- 5. Do not ingest reagent. If reagent is ingested, seek medical advice immediately.
- 6. Avoid contact with eyes. If contact occurs, flush with large quantities of water.
- 7. Follow safety precautions of the heating device used for epitope retrieval (TintoRetriever Pressure Cooker or similar).
- 8. For additional safety information refer to Safety Data Sheet for this product.
- 9. For complete recommendations for handling biological specimens, please refer to the CDC document, "Guidelines for Safe Work Practices in Human and Animal Medical Diagnostic Laboratories" (see References in this document).

Preparation of Working Solution

ImmunoDNA Digestor is a ready-to-use working solution and requires no further preparation.

Recommended Protocol

- 1. Deparaffinize sections if necessary and hydrate with distilled water.
- 2. Incubate slides with ImmunoDNA Digestor for 5 minutes at 37°C or 10 minutes at room temperature.
- 3. Rinse slides briefly in distilled or deionized water (10-20 seconds).
- 4. Proceed with IHC or ISH protocol.

Abbreviated Immunohistochemical Protocol

Abbieviatea illillationistochemicat i rotocot				
Step	ImmunoDetector AP/HRP	PolyDetector AP/HRP	PolyDetector Plus HRP	
Peroxidase/AP Blocker	5 min.	5 min.	5 min	
Primary Antibody	30-60 min.	30-60 min.	30-60 min.	
1st Step Detection	10 min.	30-45 min.	15 min.	
2nd Step Detection	10 min.	Not Applicable	15 min.	
Substrate- Chromogen	5-10 min.	5-10 min.	5-10 min.	
Counterstain / Coverslip	Varies	Varies	Varies	

Product Limitations

Due to inherent variability present in immunohistochemical procedures (including fixation time of tissues, dilution factor of antibody, retrieval method utilized and incubation time), optimal performance should be established through the use of positive and negative controls. Results should be interpreted by a qualified medical professional.

References

1. U.S. Department of Health and Human Services: Centers for Disease Control and Prevention. Guidelines for Safe Work Practices in Human and Animal Medical Diagnostic Laboratories. Supplement / Vol. 61, January 6, 2012.

Symbol Key / Légende des symboles/Erläuterung der Symbole





